arXiv:1410.8331v1 [astro-ph.SR] 30 Oct 2014

THE CoRoT DISCOVERY OF A UNIQUE TRIPLE-MODE CEPHEID IN THE GALAXY

E. Poretti[1]
INAF-Osservatorio Astronomico di Brera, Via E. Bianchi 46, 23807 Merate, Italy
ennio.poretti@brera.inaf.it
and

A. Baglin[2]
LESIA, Université Pierre et Marie Curie, Université Denis Diderot, Observatoire de Paris, F-92195 Meudon Cedex, France
and

W.W. Weiss[3]
Institute of Astronomy, University of Vienna, Türkenschanzstrasse 17, A-1180 Vienna, Austria

ABSTRACT

The exploitation of the CoRoT treasure of stars observed in the exoplanetary field allowed the detection of a unusual triple-mode Cepheid in the Milky Way, CoRoT 0223989566. The two modes with the largest amplitudes and period ratio of 0.80 are identified with the first (P_1=1.29 d) and second (P_2=1.03 d) radial overtones. The third period, which has the smallest amplitude but able to produce combination terms with the other two, is the longest one (P_3=1.89 d). The ratio of 0.68 between the first-overtone period and the third period is the unusual feature. Its identification with the fundamental radial or a nonradial mode is discussed with respect to similar cases in the Magellanic Clouds. In both cases the period triplet and the respective ratios make the star unique in our Galaxy. The distance derived from the period–luminosity relation and the galactic coordinates put CoRoT 0223989566 in the metal-rich environment of the "outer arm" of the Milky Way.

Subject headings: stars: variables: Cepheids — stars: oscillations — stars: individual (CoRoT 0223989566) — stars: interiors

1. Introduction

Double-mode Cepheids are a powerful tool to test the stellar models of supergiants, since the simultaneous excitation of two pulsation modes tightly constrains the physical parameters. In the 1990, the introduction of the new OPAL opacities (Rogers & Iglesias 1992) solved the discrepancy between the beat and pulsation masses (Moskalik et al. 1992), also reconciling the evolutionary ones (Christensen-Dalsgaard & Petersen 1995). Nowadays, the period ratio is used to investigate the metallic content of stellar systems hosting double-mode Cepheids. Sziládi et al. (2007) obtained observational evidence of the relation between the period ratios and the metal abundances by means of high-resolution spectroscopy of galactic double-mode Cepheids. The metallicity of the Large (LMC) and Small (SMC) Magellanic Clouds has been investigated by using the period ratios of hundreds of double-mode Cepheids in the framework of large-scale sur-

veys: MACHO (e.g., Alcock et al. 1999), EROS-2 (e.g., Marquette et al. 2009), OGLE-III (e.g., Soszyński et al. 2008a,b)

The involved radial modes are the fundamental (F) and the first ($1O$), second ($2O$), and third ($3O$) overtones. Typical period ratios are $1O/F$=0.71, $2O/1O$=0.80, $3O/1O$=0.68. However, the large-scale surveys revealed other particular subclasses (for a review see Moskalik 2014): stars with a nonradial mode in close proximity to the dominant $1O$ mode and double-mode pulsators with a period ratio of 0.60-0.64.

On the other hand, triple-mode Cepheids are extremely rare. The current statistics (Moskalik 2014) report six cases of $1O/2O/3O$ stars (three in the LMC, one in the SMC, and two in the galactic bulge) and four cases of $F/1O/2O$ stars (two in the LMC, two in the SMC). The two triple-mode Cepheids in the galactic bulge have extremely short periods: the $1O$ periods are 0.295 d and 0.230 d, corresponding to F periods shorter than 0.4 d. The modeling of three periods and the evolutionary calculations are expected to place strong constraints both on Cepheid mass–luminosity relations and on the internal physics.

The variability of GSC 0746-01186 was discovered in the ASAS survey (ASAS 064135+0756.6; Pojmanski 2002). The star was classified as a Cepheid with P=1.28859 d, with an amplitude of 0.41 mag and $< V >$=12.48. Later, Khruslov (2009) identified it as a double-mode Cepheid with periods P_1=1.28861 d and P_2=1.03153 d. These periods were confirmed by the analysis of the NSVS data (Woźniak et al. 2004). The ratio of 0.8005 suggested a pulsation in the $1O$ and $2O$ modes. GSC 0746-01186 ≡ ASAS 064135+0756.6 ≡ SRa01b.16229 was re-observed during the BEST II survey (Berliner Exoplanet Search Telescope; Klagivik et al. 2013). The double-mode pulsation and the period ratio were both confirmed. The space mission CoRoT (Baglin et al. 2006) monitored the star in a serendipitous mode since it is located in a field contiguous to that of the open cluster NGC 2264, the main target of the first short run in the anticenter direction (SRa01). We definitely cross-identify CoRoT 0223989566≡GSC 0746-01186≡ASAS 064135+0756.6≡2MASS 06413457 +0756396 when searching for Cepheids in the CoRoT data base (Poretti et al., in preparation).

2. The analysis of CoRoT 0223989566 data

CoRoT 0223989566 was observed continuously for 23.4 d from 2008 March 7 to March 31. The EXODAT catalog (Deleuil et al. 2009) reports a spectral type of A5 III, an E_{B-V}=1.1 mag (both from SED analysis), and a very low contamination level due to close stars, i.e., 0.004 in a range from 0 to 1. The "CoRoT Variability Classifier" (CVC) automated supervised method (Debosscher et al. 2009) correctly suggests a classical or a double-mode Cepheid. The measurements of CoRoT 0223989566 were performed in the 512-s mode from JD 2454533.4120 to 2454536.008 and then in the 32-s mode until the end of the observations, at JD 2454556.795.

All the CoRoT measurements were taken into account for the initial frequency analysis. The outliers were removed by means of a cross-check between the flags provided by the reduction pipeline and a visual inspection of the light curve. Outliers are for most measurements that suffer from hot pixels during the passage through the South Atlantic Anomaly. The final time series is composed of 52,148 measurements: they were obtained in the chromatic mode, but here we discuss the white measurements only since we are interested in a detailed frequency analysis requiring a high signal-to-noise ratio (S/N). The high-precision, high duty-cycle CoRoT photometry provides us with an excellent representation of the beating between two commensurable periods (Fig. 1, top panel). The light curve is a succession of two "bright" maxima followed by two "faint" ones. The four maxima span the short beating period of 5.16 d=$4P_1$=$5P_2$.

The frequency analysis was performed by means of the iterative sine-wave, least-squares fitting method (Vaniĉek 1971). It was applied to boththe original CoRoT time series composed of 32-s and 512-s exposures and to a new data set obtained by grouping 16 consecutive 32-s measurements into a single 512-s point. The frequency values were refined by the MTRAP algorithm (Carpino et al. 1987), allowing us to search for the best fit by keeping the values of the harmonics and the combination terms locked to the independent frequencies. The analysis of the CoRoT data identified the two frequencies corresponding to the

periods already known (Fig. 2, the power spectra from the original time series are shown). The peak at f_1=0.776 d^{-1} is by far the highest in the power spectrum, but $2f_1$ and f_2=0.970 d^{-1} are also immediately noticeable (panel *a*). The f_2 peak and the combination terms f_1+f_2 and f_2-f_1 were detected after prewhitening the data with f_1 and its harmonics (panel *b*). The four terms f_1, f_2, f_1+f_2, and f_2-f_1 are the most common in the solutions of the light curves of double-mode Cepheids obtained from ground-based data (Pardo & Poretti 1996). We expected that the subsequent analysis of the CoRoT time series would disclose the rich ensemble of combination terms between the harmonics of f_1 and f_2. Therefore, it was a great surprise to detect a new independent term f_3=0.529 d^{-1}, at the next step (panel *c*). The presence of the f_1+f_3 peak was particularly important since it immediately ruled out that f_3 was due to the variability of a close, unresolved star in the CoRoT mask or to a binary companion. After this, the combination terms between the harmonics of f_1, f_2, and f_3 were detected step-by-step, as shown in panel *d* for a subset of nine of them. The final solution of the CoRoT light curve was provided by the independent terms f_1=0.776006 $\pm$ 0.000002 d^{-1}, f_2=0.969787 $\pm$ 0.000016 d^{-1}, and f_3=0.529632 $\pm$ 0.000143 d^{-1}, their harmonics (up to $7f_1$, $2f_2$, and $2f_3$) and a set of 23 other combination terms, for a total of 34 components. Table 1 lists the ordered structure of the combination terms and the least-squares solution obtained from the time series composed of the 512-s and 32-s measurements, which leaves a residual r.m.s. of 0.0010 mag. The error bars on the amplitudes $\sigma(A) = 6 \cdot 10^{-6}$ mag and on the phases $\sigma(\phi_i) = \sigma(A)/A_i$ can be immediately derived following Montgomery & O'Donoghue (1999).

The discovery of f_3 opened the possibility that other modes were excited and we performed an additional analysis to find them. The power spectra of the entire data set show some peaks close to f_1 and, much less relevant, to f_2. The highest one was at f=0.728 d^{-1} with an amplitude 0.008 times that of f_1. The nature of these peaks is ambiguous. They could have a stellar origin and be due to a long-term modulation of the main oscillations, observed in some $1O/2O$ Cepheids (Moskalik 2014). However, we cannot rule out the possibility that this long-term effect is actually due to satellite drift and/or detector aging. The analysis of the light curve of the residuals helped us to clarify this point. After the prewhitening with 34 components the data show residual oscillations, up to $\pm$2 mmag (Fig. 1, middle panel). The amplitudes of the peaks found in the frequency analysis of the residuals are smaller than 0.1 mmag. An erratic nature could be more plausible, but not fully convincing. Therefore, we calculated a new set of residuals by subdividing the 23.4-d time baseline into six contiguous subsets spanning 3.9 d each. The frequency values were kept fixed, but the amplitudes and phases were recalculated for each subset. Then, the residuals obtained from these subsets were merged. The residual light curve thus obtained is almost flat and the erratic oscillations have completely disappeared (Fig. 1, bottom panel). We can infer that the erratic oscillations are due to instrumental effects that are affecting the CoRoT photometry in such a way that the usual technique of prewhitening was not able to clean (see also the case of CoRoT 101155310; Poretti et al. 2011). The frequency analysis of the residuals after subtracting the 34 components from the six subsets (Fig. 2, panel *e*) did not detect any peak below 1.0 d^{-1}, where we expected the independent modes we were looking for.

Therefore, we can conclude that after f_3 no other independent mode is detectable in the light curve of CoRoT 0223989566. The noise level of the power spectrum of the residuals (Fig. 2, (panel *e*) is around 0.004 mmag in the 0-3 d^{-1} region and increases to 0.012 mmag in the 5.0-7.0 d^{-1} region. Some of the peaks visible in the 5.0-7.0 d^{-1} region are close to combination terms (e.g., $6f_1+f_3$=5.18 d^{-1}, $7f_1$=5.43 d^{-1}, $7f_1+f_2$=6.40 d^{-1}, $6f_1+2f_2$= 6.59 d^{-1}), but their amplitudes (0.028, 0.040, 0.028, and 0.030 mmag, respectively) are below the heuristically accepted threshold of S/N=3.5 (Breger et al. 1993).

For the sake of completness, we successfully verified that the f_3 component was also detectable in the CoRoT data before removing the outliers and that the solution of the time series composed of the grouped 512-s points supplies the same combination terms of the solution listed in Table 1.

3. Discussion

The analysis of the data of CoRoT 0223989566 returned three independent periods: P_1=1.2886 d, P_2=1.031140 d, and P_3=1.888538 d. This triplet is completely new among galactic Cepheids. The period ratios are P_1/P_2=0.8002, P_2/P_3=0.546, and P_1/P_3=0.682.

3.1. The Petersen Diagram

Figure 3 shows the Petersen diagram of the galactic double-mode Cepheids. There is little doubt on the fact that P_1 and P_2 can be typified with the $1O$ and $2O$ modes, as is done in the galactic, LMC, and SMC double-mode Cepheids. P_3 is giving us more trouble. The ratio 0.68 has been observed in two double-mode LMC Cepheids (Soszyński et al. 2008a) and in six triple-mode Cepheids ($1O/2O/3O$) in both the Magellanic Clouds (Moskalik 2014). The related periods were identified with those of the $3O$ and $1O$ modes. However, this is not the case for CoRoT 0223989566: since P_3 is longer than P_1, it should be the $1O$ mode and P_1 the $3O$ one, contradicting the previous robust identification of P_1 as the $1O$ mode.

The fact that P_3 is much longer than P_1 and P_2 naturally suggests its identification with the F radial mode. However, the corresponding ratio $1O/F$=0.682 is lower than the usual one (0.694-0.746; Moskalik 2014). Moreover, Fig. 3 shows how the $1O/F$ ratio is expected to increase toward short F-periods, thus making the 0.682 ratio of CoRoT 0223989566 still more peculiar. There is only one example of a ratio of 0.68 typified as a $1O/F$ one, i.e., J045917-691418 (P_0=3.08 d and P_1=2.10 d) in the LMC (it is the apparently lowest outlier in Fig. 3 of Marquette et al. 2009). The unusual ratio is explained in terms of an high-metallicity of $Z = 0.030$ (see also Fig. 3 in Buchler & Szabó 2007), while the mean metallicity of LMC double-mode Cepheids is 0.004 [1]. The fundamental period of CoRoT 0223989566 is shorter than that of J045917-691418 and a still higher metallicity is necessary to include the $(\log P, P_1/P_0)$ location of CoRoT 0223989566 between the limits where F and $1O$ are both unstable (Buchler & Szabó 2007).

Therefore, if P_3 is actually the F mode, CoRoT 0223989566 is a very particular star: not only the unique $F/1O/2O$ Cepheid in the Galaxy but also one with an unusual $1O/F$ ratio. In any case, CoRoT 0223989566 is the triple-mode Cepheid showing the longest periods, both in the Galaxy and in the Magellanic Clouds. In the context of the galactic Cepheids, it should be noted that P=1.8 d seems to be the shortest F period among the $1O/F$ double-mode stars and the longest $1O$ period among the $2O/1O$ ones (Fig. 3).

3.2. The Analysis of the Fourier Parameters

The Fourier decomposition could be used to disentangle the matter, since the first harmonics were found for the three periods of CoRoT 0223989566. We calculated the Fourier parameters $R_{ij} = A_i/A_j$ and $\phi_{ij} = j\ \phi_i - i\ \phi_j$ from the amplitude (A_i) and phase (ϕ_i) coefficients of the i and j harmonics of the three independent frequencies f_1, f_2, and f_3.

The parameters ϕ_{21}=4.1480±0.0006 rad and R_{21}=0.2108 ± 0.0001 of f_1 are in excellent agreement with those of the $1O$ modes of the $1O/2O$ Cepheids. In a similar way, the Fourier parameters of f_2 (ϕ_{21}=5.69±0.05 rad, R_{21}=0.014±0.001) are exactly located among those of the $2O$ modes (see Figs. 2 and 3 in Alcock et al. 1999, for the LMC Cepheids). We conclude that CoRoT 0223989566 does not show any particularity as a $1O/2O$ double-mode pulsator.

The situation is slightly different when we consider f_3. The ϕ_{21}=4.05±0.12 rad value observed for f_3 is on the extension of the F-mode progression (see Fig. 4 in Pardo & Poretti 1996), but R_{21}=0.047±0.005 is a small value for the F-mode, usually greater than 0.20. V371 Per is another double-mode Cepheid where the R_{21}=0.15 of the F-mode is below the 0.20 limit and smaller than the R_{21}=0.22 of the $1O$-mode (Wils et al. 2010). We can suppose that the small amplitude of the F-mode oscillation reduces the R_{21} value. It is also noteworthy that V371 Per shows an unusual high $1O/F$ period-ratio (0.731), explained in terms of a

[1] We cannot rule out that J045917-691418 is actually a $3O/1O$ pulsator since the period ratio agrees very well with those of the double-mode and triple-mode Cepheids recalled above. We verified that a F-period of 3.08/0.715=4.31 d matches the $P - L$ relation as well as 3.08 d.

metal deficiency ($-1 <$[Fe/H]< -0.7). Therefore, it seems that given stars could deviate from the most common ratios.

Another possibility is to identify f_3 as a nonradial mode. Nonradial modes have been detected in $1O$ LMC Cepheids and $F/1O$ double-mode LMC Cepheids, always closely to the $1O$ mode ($\Delta f < 0.13$ d^{-1}; Moskalik 2014). If we suppose a canonical $F/1O$ frequency ratio of 0.72, we have $F = 1O \cdot 0.72$=0.559 d^{-1}. The difference $0.559 - 0.529 = 0.030$ d^{-1} is still in the range where we can think of a resonance. However, also under these assumptions CoRoT 0223989566 remains an unique and challenging case since we have to model the excitation of a nonradial mode close to the expected – but not observed – F radial mode.

3.3. The Location of CoRoT 0223989566 in the Milky Way

The $P - L$ relation for galactic Cepheids $M_V = -2.999(\pm0.097) \log P - 0.995(\pm0.112)$ (Ngeow & Kanbur 2004) supplies similar M_V values if we consider f_3=0.529 d^{-1} or $0.776 \cdot 0.72 =$ 0.559 d^{-1} as the frequency of the F radial mode, i.e., $M_V = -1.82 \pm 0.12$ and $M_V = -1.75 \pm$ 0.12, respectively. The color excess E_{B-V}=1.1 (EXODAT; Deleuil et al. 2009) yields $A_V =$ 3.2 E_{B-V}=3.5, R=3.2 from Tammann et al. (2003). The distance modulus $V - M_V = 12.5 + 1.8 - 3.7 =$ 10.8 and the galactic coordinates (l=204.°7407, b=+01.°4386) put CoRoT 0223989566 at 1.4 kpc from the Sun, behind and farther than NGC 2264 (9.3; 202.°936 +02.°196). The large color excess seems confirmed by available photometry: V=12.50 (Pojmanski 2002) and B=13.6 (GSC 2.2 catalog). Smaller values close to E_{B-V}=0.4 have been measured in that direction (Jenkins & Tripp 2011). Such a value would push CoRoT 0223989566 at 4.0 kpc from the Sun. In any case, the resulting galactocentric distances R_G=9.8 kpc (E_{B-V}=1.1) or R_G=12.2 kpc (E_{B-V}=0.4) or any intermediate one obtained from $0.4 < E_{B-V} < 1.1$ mag) place CoRoT 0223989566 on the extreme "outer arm", not far from the B0 II star HD 43818, i.e., in an environment known to be metal rich (Tripp & Song 2012).

Therefore, the very high metallicity could cause the $1O/F$ period ratio to decrease in the opposite sense of what the low metallicity probably does for V371 Per (Wils et al. 2010). The fact that the $2O/1O$ ratio is normal is not surprising, since the metallicity affects mainly the $1O/F$ ratio (Beaulieu et al. 1997; Marquette et al. 2009). Other evolutionary effects linked with the particular galactic location, as the first crossing of the instability strip of a young object or an anomalous mass with respect to other double-mode Cepheids, can contribute to the unusual period ratio.

4. Conclusions

The intensive CoRoT monitoring of CoRoT 0223989566 discovered a unique case among triple-mode galactic Cepheids. The 0.682 ratio between the two longest periods is quite unusual. If interpreted as a $1O/F$ ratio, the fact that CoRoT 0223989566 belongs to a metal-rich environment like the "outer arm" of the Milky Way could explain the value. We also note that the periods of CoRoT 0223989566 are much longer than those of the triple-mode Cepheids detected in the galactic bulge. If not a radial mode, the excitation of an isolated nonradial mode with a period much longer than that of the $1O$ mode is also unusual for high-amplitude pulsators. As expected (e.g., Moskalik & Dziembovski 2005; Moskalik 2014), the discovery of a new member of the rare class of triple-mode Cepheids set new observational constraints on the stellar parameters of these variables and, in a more wide context, on their evolutionary models.

The CoRoT space mission has been developed and operated by CNES, with contributions from Austria, Belgium, Brazil, ESA (RSSD and Science Program), Germany, and Spain. This research has made use of the ExoDat Database, operated at LAM-OAMP, Marseille, France, on behalf of the CoRoT/Exoplanet program. The present study has used the SIMBAD data base operated at the Centre de Données Astronomiques (Strasbourg, France).

REFERENCES

Alcock, C., Allsman, R.A., Alves, D. et al. 1999, ApJ, 511, 185

Baglin, A., et al., 2006, in "The CoRoT Mission, Pre-Launch Status, Stellar Seismology and

Planet Finding", ed. M. Fridlund, A. Baglin, J. Lochard, L. Conroy (ESA SP-1306, Nordwijk, ESA), 33

Beaulieu, J.-P., Krockenberger, M., Sasselov, D.D., et al. 1997, A&A, 321, L5

Breger, M., Stich, J., Garrido, R., et al. 1993, A&A, 271, 482

Buchler, J.R., & Szabó, R. 2007, ApJ, 660, 723

Carpino, M., Milani, A., & Nobili, A.M. 1987, A&A, 181, 182

Christensen-Dalsgaard, J., & Petersen, O.J. 1995, A&A, 299, L17

Debosscher, J., Sarro, L. M., López, et al., A&A, 2009, 506, 519

Deleuil, M., Meunier, J. C., Moutou, et al. 2009, AJ, 138, 649

Jenkins, E.B., & Tripp, J.M. 2011, ApJ, 734, 65

Khruslov, A.V. 2009, Per. Zvezdy, 9, 31

Klagivik, P., Csizmadia, Sz., Pasternacki, T., Fruth, T., Erikson, A., et al. 2013, ApJ, 773, 54

Marquette, J.B., Beaulieu, J.P., Buchler, J.R., et al. 2009, A&A, 495, 249

Montgomery, M.H., & O'Donoghue, D. 1999, Delta Scuti Star Newsletter, 13, 28

Moskalik, P. 2014, in IAU Symp. 301 "Precision Asteroseismology", ed. J.A. Guzik, W.J. Chaplin, G. Handler, & A. Pigulski, (Cambridge: Cambridge Univ. Press), 249 (arXiv:1401.7271)

Moskalik, P., Buchler, J.R., & Marom, A. 1992, ApJ, 385, 685

Moskalik, P., & Dziembovski, W.A. 2005, A&A, 434, 1077

Ngeow, C.-C, & Kanbur, S.M. 2004, MNRAS, 349, 1130

Pardo, I. & Poretti, E., 1996, A&A, 324, 121

Pojmanski, G. 2002, Acta Astron., 52, 397

Poretti E., Rainer M., Weiss, W.W., et al. 2011, A&A, 528, A147

Rogers, F.J, Iglesias, C.A. 1994, Science, 263, 50

Soszyński, I., Poleski, R., Udalski, A., et al. 2008a, Acta Astron., 58, 153

Soszyński, I., Poleski, R., Udalski, A., et al. 2008b, Acta Astron., 58, 163

Sziládi, K., Vinkó, J. Poretti, E., Szabados, L., & Kun M. 2007, A&A, 473, 579

Tammann, G.A., Sandage, A., & Reindl, B. 2003, A&A, 404, 423

Tripp, T.M., & Song, L. 2012, ApJ, 746, 173

Vaniĉek, P. 1971, Ap&SS, 12, 10

Wils, P., Henden, A.A., Kleidis, S., Schmidt, E.G., & Welch, D.L. 2010, MNRAS, 402, 1156

Woźniak, P.R., Vestrand, W.T., Akerlof, C.W., et al. 2004, AJ, 127, 2436

This 2-column preprint was prepared with the AAS LaTeX macros v5.2.



Fig. 1.— *Top panel:* CoRoT photometry (white light) of CoRoT 0223989566. *Middle panel:* residuals from a solution calculated on the entire data set spanning 23.4 d. *Bottom panel:* residuals from the solutions of six subsets spanning each 3.9 d. Gray points: 32-s exposures; black points: 512-s exposures or 512-s averages.



Fig. 2.— Frequency analysis of the time series composed of the 512-s and 32-s measurements after subsequent prewhitenings. Panels *a* and *b*: detection of f_1, f_2, and combination terms. Panel *c*: detection of f_3 and of the combination term f_1+f_3. Panel *d*: detection of a subset of the combination terms between f_1, f_2, and f_3. Panel *e*: power spectra of the residuals obtained from six subsets.



Fig. 3.— Petersen diagram of galactic double-mode Cepheids (open circles, F and $1O$ stars; open squares, $1O$ and $2O$ pulsators). The triple-mode Cepheid CoRoT 0223989566 is indicated with filled symbols.

Table 1: The identification of the 34 significant frequencies detected in the data of CoRoT 0223989566. The amplitudes and phases obtained from the original 32-s and 512-s measurements are listed.

ID	Frequency [d^{-1}]	Amplitude [mag]	Phase [0,2π]
f_1	0.776006	0.071792	0.6504
f_2	0.969787	0.009816	3.8558
f_3	0.529632	0.001181	1.5836
$2f_1$	1.552012	0.015132	5.4488
$3f_1$	2.328018	0.003827	3.8502
$4f_1$	3.104024	0.000723	2.4672
$5f_1$	3.880030	0.000198	0.1619
$6f_1$	4.656036	0.000089	4.7833
$2f_2$	1.939574	0.000137	0.8405
$2f_3$	1.059264	0.000056	0.9304
f_2–f_1	0.193781	0.002828	1.0666
f_1+f_2	1.745793	0.003041	2.9196
$2f_1$+f_2	2.521799	0.001057	1.7247
$3f_1$+f_2	3.297805	0.000160	2.1453
$4f_1$+f_2	4.073811	0.000116	1.6367
$5f_1$+f_2	4.849817	0.000101	0.3820
$6f_1$+f_2	5.625823	0.000066	5.8167
$2f_1$–f_2	0.582225	0.000324	5.1539
$3f_1$–f_2	1.358231	0.000066	0.6502
$4f_1$–f_2	2.134237	0.000056	3.2205
f_1+$2f_2$	2.715580	0.000162	6.0254
$2f_1$+$2f_2$	3.491586	0.000104	5.5258
$3f_1$+$2f_2$	4.267592	0.000104	4.8089
$4f_1$+$2f_2$	5.043598	0.000080	3.8391
$5f_1$+$2f_2$	5.819604	0.000055	2.5362
$2f_2$–$2f_1$	0.387562	0.000056	1.5524
$4f_1$+f_3	3.633656	0.000061	0.9406
f_1+f_3	1.305638	0.000699	6.0973
$2f_1$+f_3	2.081644	0.000286	4.4819
$3f_1$+f_3	2.857650	0.000153	2.8912
f_1–f_3	0.246374	0.000271	4.3864
f_2+f_3	1.499419	0.000192	2.9091
f_2–f_3	0.440155	0.000051	6.0342
f_1+f_2+f_3	2.275425	0.000069	1.9800
N measurements: 52148			
T_0= HJD 2454546.3070			
Residual r.m.s. 0.00104 mag			